

NATURALLY
DERIVED



MITE TONIC

A nutritional supplement, added to the drinking water of laying hens, that helps improve the natural resistance of poultry through natural plant extracts.

5L



APPLICATION TO POULTRY
DRINKING WATER



DILUTION
500mL / 1000L

COMPLEMENTARY FEED FOR LAYERS

MiteTonic contributes to improve the natural resistance of layer hens in their environment.

DIRECTIONS FOR USE: MiteTonic must be administered in drinking water at the dosage of 0,5 L for 1000 L of water, at the rate of 200 ml per hen.

Protocol 1. Continuous application from flock introduction:
1 day per week, during 6 weeks. Renew this treatment every month for 2 consecutive days.

Protocol 2. Intermediate application in case of deficiencies:
For 4 consecutive days plus 2 days three weeks later and 2 days four weeks later. Renew this treatment every 15 to 20 weeks for 4 consecutive days.

ANALYTICAL CONSTITUENTS

HUMIDITY.....	92.69
CRUDE FAT	0.00
CRUDE ASH	1.56
LYSINE	0.00
CRUDE PROTEIN.....	0.00
CRUDE FIBRE.....	0.00
SODIUM	0.07

METHIONINE 0.0

STORAGE OF PRODUCT

Store in a cool and dry place, keep at ideal temperature between 5°C and 25°C, away from direct light and heat sources. MiteTonic should not be stored near electrical devices.

COMPOSITION PER L

Purified Water, Magnesium chloride, Sodium chloride.

ADDITIVES:

3a - Vitamins, provitamins and substances chemically defined:

Vitamin C (ascorbic acid) 12 380 mg/L

2b - Aromatic substances:

Litsea cubeba 3000 mg Thymus vulgaris 1670 mg
Origanum vulgare 1670 mg Inula helenium 1670 mg.

WITHDRAWAL PERIOD:

None

LODI UK LIMITED

Building 69, Third Avenue, Pensnett Trading Estate,
Kingswinford, West Midlands DY6 7FD
Tel: 01384 404242
Email: sales@lodi-uk.com
www.lodi-uk.com

Batch Nr. / Exp. date :

GMP021417 Armosa BV
NU-CHI-02007 PA-LAB-03235



5L