

TEAR HERE

RESEALABLE STRIP



SAPPHIRE Grain 25



Lodi's Gems

Rat and Mouse KILLER

SINGLE FEED

150g Pouch

150g



Special labelling provisions for Ireland: Use biohazard safety and sustainability. It is illegal to use this product for uses or in a manner other than prescribed on this label. For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-809-2166), retain the label for reference. Where control of rodents has not been achieved by a control program after 35 days, the user should seek advice from the product supplier or call a pest control service. Do not use this product for permanent or pulse-baiting.

Search for and remove dead rodents during treatment, at least as often as baiting stations are inspected. Dispose of dead rodents in accordance with local requirements. In the UK, poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or collected by a specialist contractor or the local authority. Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait. Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing of them. This product contains a blinding agent and a dye.

Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment. This product contains an anticoagulant substance. If ingested, symptoms, which may include nosebleed and bleeding gums. Remove rodents, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only. In case of: Dermal exposure, wash skin with water and then with soap. Eye exposure, rinse eyes with first-aid liquid or water. Keep eyes lids open at least 10 minutes. EYE EXPOSURE, rinse eyes with water. Never give anything by mouth to unconscious persons. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice. Contact a veterinary surgeon in case of ingestion by a pet. Bait stations must be labelled with the following information: "do not move or open", "contains a rodenticide", "product name or authorisation number", "active substance(s)" and "in case of accident, call a poison centre". Hazardous to wildlife.

Instructions for safe disposal of the product and its packaging: UK: At the end of the treatment, dispose unopened bait and the packaging in accordance with EPA requirements for the disposal of hazardous waste. Use of gloves is recommended. In the UK bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or taken to a civic amenity site. For information on safe disposal contact the local authority. At the end of the treatment, remove potential food and drinking water (plug holes, remove potential rodenticide bait or the bait stations at the end of the treatment period. Remove the remaining bait or the bait stations and directly exposed skin after using the product. Do not eat, drink or smoke. Wash hands and directly exposed skin after using the stations near water drainage systems where they can come into contact with water. When using farm animals and other non-target animals. Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Do not place bait open the sachets containing the bait. Place bait stations out of the reach of children, birds, pets, vicinity where rodenticide activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). Where possible, bait stations must be fixed to the ground or other structures. Do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. Bait stations should be placed in the immediate vicinity where rodenticide activity has been observed (e.g. traps) should be considered. Remove accompanying the product or provided at the point of sale before using it. Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered. Remove insects, symptoms, which may include nosebleed and bleeding gums. Remove rodents, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only. In case of: Dermal exposure, wash skin with water and then with soap. Eye exposure, rinse eyes with first-aid liquid or water. Keep eyes lids open at least 10 minutes. EYE EXPOSURE, rinse eyes with water. Never give anything by mouth to unconscious persons. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice. Contact a veterinary surgeon in case of ingestion by a pet. Bait stations must be labelled with the following information: "do not move or open", "contains a rodenticide", "product name or authorisation number", "active substance(s)" and "in case of accident, call a poison centre". Hazardous to wildlife.

Conditions of storage and shelf-life of the product under normal conditions of storage: Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals. Shelf life: 2 years.

Use-specific instructions for use: Mice: The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards. Re-visit the bait stations if necessary. Re-fill bait when necessary. Re-fill bait when necessary. Re-visit the bait stations in areas non-labile to flooding. Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt. The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait stations are intact and to remove rodent bodies. Application rate(s) and frequency: For rats: 45 - 60 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be 5 - 10 m apart. For mice: 10 - 25 g of bait per bait station. If more than 1 bait station is needed, the minimum distance between bait stations should be 2 - 5 m apart.

Target organisms: Rattus norvegicus Brown Rat; Rattus rattus Roof Rat; House Rat, Mus musculus House mouse. Application method: Ready-to-use bait to be used in tamper-resistant bait stations. Application method: Ready-to-use bait to be used in tamper-resistant bait stations. Tamper-resistant bait stations: Strong enough to prevent entry or destruction by dogs. Strong enough to prevent entry or destruction by children under six years of age using hands, feet or objects, toys etc.). Lockable or sealable so that children and dogs cannot gain access through the opening or mechanism used to fill the bait compartments. Resistant to destruction or weakening from exposure to typical non-catastrophic weather (e.g. direct sunlight, extremes of temperature and humidity, rain, snow etc), use for mice and rats / outdoor around buildings for rats only. Field of use: Indoor use for mice and rats / outdoor around buildings for rats only. User category: For general public use. Authorisation No Ireland: IE/BPA-70677-001 Authorisation No Great Britain: GB-2016-1056-0003 NI-2016-1056-0003

SAPPHIRE GRAIN 25 - Brodifacoum (CAS 56073-10-0) : 0.0025% Warning: (H373) May cause damage to organs (H403) through prolonged or repeated exposure. P102 - Keep out of reach of children. P220 - Keep/store away from food, drink and animal feedstuffs. P262 - Avoid breathing dust. P282 - Do not get on skin. P270 - Do not eat, drink or smoke when using this product. P301+310 - IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician. P314: Get medical advice/attention if you feel unwell. P404+405 - Store locked up in a closed container. P501 Dispose of contents/container in accordance with applicable regulations.

Authorisation holder (GB): LODI UK, Unit 104, Potter Space, Z Kidderminster Road, Cutnall Green, Droitwich, WR9 0NS Grand Fougeray, 35390 France

RESEALABLE STRIP