

# INSECTICIDE FOR PROFESSIONAL USE AGAINST FLYING AND CRAWLING INSECTS SUCH AS FLIES AND RED MITES.

FOR USE IN AND AROUND FARM BUILDINGS, DAIRIES, MILKING PARLOURS, ANIMAL REARING FACILITIES, ANIMAL HOUSING INCLUDING POULTRY HOUSES (SURFACE SPRAY ONLY), ANIMAL TRANSPORTATION AND SLAUGHTERHOUSES AND WASTE MANAGEMENT SITES (INDOORS) INCLUDING WASTE TRANSFER STATIONS AND MBT TREATMENT PLANTS, REFUSE DISPOSAL AND WASTE STORAGE AREAS (INCLUDING REFUSE TIPS AND LANDFILL SITES).

**APPLICATION METHOD:** Surface spray

To avoid risks to human or animal health, or the environment, comply with the instructions for use.

## STATUTORY CONDITIONS OF APPROVAL RELATING TO USE

**FOR USE ONLY AS AN INSECTICIDE  
FOR USE ONLY BY PROFESSIONAL  
OPERATORS.**

**APPLICATION RATE:** Apply 1 litre of product per 20 square metres of surface or spray short bursts directly at insects.

Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended) and the Control of Pesticides Regulations (Northern Ireland) 1987 (as amended), including complying with all Conditions of Approval.

**READ ALL PRECAUTIONS BEFORE USE**  
**HSE No: 11103-011-GB, 11103-011-NI**

## RISK MITIGATION MEASURES:

- \*The COSHH (Control of Substances Hazardous to Health) Regulations 2002 (as amended) apply to the use of this product at work
- \*HAZARDOUS TO BEES.
- \*All bats are protected under The Wildlife and Countryside Act 1981. Before treating any structure used by bats, consult Natural England, NatureScot, Natural Resources Wales or the Northern Ireland Environment Agency
- \*Not to be used on pet bedding, pet sleeping quarters or where pets frequently lie.
- \*Do not apply directly to animals.
- \*Avoid excessive contamination of coveralls.
- \*Do not contaminate foodstuffs, eating utensils or food contact surfaces.
- \*Action should be taken to prevent foraging bees gaining access to treated nests, preferably by removing the combs or blocking the nest entrance.
- \*Protect milk machinery and milk containers from contamination.
- \*Exclude all persons and animals during treatment.
- \*Wash hands and exposed skin before meals and after use.
- \*Cover all water storage tanks before application.
- \*This material and its container must be disposed of in a safe way.
- \*Do not apply to clothing and bedding.
- \*Do not touch treated surfaces until dry
- \*Keep in a safe place.
- \*Collect eggs and remove feed, exposed water and milk before application.

## DIGRAIN ULTRA DAIRY FLY SPRAY

### ACTIVE INGREDIENTS:

Cypermethrin (CAS 52315-07-8) 0.102% w/w  
Tetramethrin (CAS 7696-12-0) 0.209% w/w



### WARNING

- H410 Very toxic to aquatic life with long lasting effects.
- EUH208 Contains 2-METHYLISOTHIAZOL-3(2H)-ONE. May produce an allergic reaction.
- EUH208 Contains Benzyl Alcohol. May product an allergic reaction.
- P273 Avoid release to the environment.
- P391 Collect spillage
- P501 Dispose of contents/container according to the regulation.

Lodi UK Limited, Unit 104, Potter Space,  
7 Kidderminster Road, Cutnall Green,  
Droitwich, WR9 0NS  
Tel: 01384 404242

**UFI: ARQ9-H88J-C70H-FYUV**

### MARKETING COMPANY:

Lodi UK Limited, Unit 104, Potter Space,  
7 Kidderminster Road, Cutnall Green, Droitwich,  
WR9 0NS  
Tel: 01384 404242  
E-mail: sales@lodi-uk.com  
www.lodi-uk.com

### EU ADDRESS:

LODI S.A.S, Parc d'Activites des Quatre Routes,  
35390 Grand-Fougeray, FRANCE

**DIGRAIN**

**ULTRA  
DAIRY  
FLY  
SPRAY  
RTU**



 **Ultra knockdown against flies & crawling insects**

 **For use in dairies, animal housing & rearing facilities**

**5 Litres**

**FOR PROFESSIONAL USE**

  
**LodiUK**