

IMIDASECTANTS



6x35g

▶ GEL BAIT INSECTICIDE AGAINST ANTS

For use by Professionals only

Formulation: Ready-to-use gel

Active substance: Imidacloprid

Chemical family: Nitroguanidines (Neonicotinoids)

Concentration: 0.01% w/w

Authorisation Number: GB-2016-1043

Authorisation holder:

Sharda Agrochem Limited
201, Cervantes House, 5-9, Headstone Road,
Harrow, Middlesex, HA1 1PD

Distributor:

LODI UK, Unit 104, Potter Space,
7 Kidderminster Road, Cutnall Green,
Droitwich WR9 0NS

Tel: 01384 404242

www.lodi-uk.com

sales@lodi-uk.com

Emergency phone: UK +44 (0) 1865 407333

IMPORTANT: USE INSECTICIDES WITH PRECAUTION. BEFORE USE, READ THE LABEL.

Respect the uses, doses, conditions and precautions on the packaging which have been determined based on product characteristics and applications for which it is recommended.

The treatment should be performed according to good practice taking into account under your responsibility all the specific factors that may be involved and are beyond our control (materials used, application conditions, product mix, etc.)

The manufacturer guarantees the quality of its products sold in their original packaging.



Hazard Statement:

H411 Toxic to aquatic life with long lasting effects.

Precautionary statements:

P273 Avoid release to the environment.

P391 Collect spillage.

P501 Dispose content/containers according to national legislation.

Observe the instructions to avoid risks to man and the environment.

Emergency phone: UK +44 (0) 1865 407333

Size: 6x35 g

Batch No.:

Manufacture date:

Expiry date: 4 years from manufacturing date



UFI: PEDV-G68N-F904-U6NY



8 904150 079836

Target insects:

Pharaoh ants (*Monomorium pharaonis*), Argentine ants, (*Linepithema humile*), Black garden ants (*Lasius niger*)

Mode of action:

The active ingredient imidacloprid acts by binding to the insect's neurons. This binding causes a disturbance in the transmission of nerve impulses which is lethal to the target insect.

Ants are attracted by nutritional ingredients in the bait. They spread the gel by moving and contaminate their peers, causing poisoning by contact and ingestion regardless of their stage of development (eggs, larvae, nymphs, pupae, imagines, adults).

Use instructions: Before treatment, remove as much as possible all natural source of food for ants (waste, food scraps ...) from the infested area to encourage the ingestion of the gel.

Place the gel in drops or thin lines near ant trails, points of entry and their nests.

Where presence of ants is only suspected or detected sporadically, it is recommended to use the gel in a bait tray. Make inspections once a week, check the treated area and if necessary, replace the gel.

Do not apply the product on absorbing surfaces.

Apply the product away from direct sunlight or heat sources (e.g. do not place it under a radiator).

Avoid continuous use of the product.

Inform the distributor if the treatment is ineffective.

The outdoor application should only be conducted on paved surfaces (do not apply on bare soil).

IMIDASECT ANTS is recommended for use in:

Domestic premises (including kitchens)

Food handling places: Processing: Food manufactures, kitchens, slaughterhouses etc. Storage: Food retailers, warehouses, raw material stores, silos (except where grain is stored) etc. Preparation: Restaurants, commercial kitchens, etc.

Public buildings: Continuously occupied areas including hospital wards, hotels, public baths, municipal buildings, churches, halls, community centres, cinemas, etc.

Application dose:Indoor use:

- 1-2 drops of gel per m²

Apply drops of gel having a diameter of about 1 cm (approximately 0.2 g)

Outdoor use:

- 1 drop of gel directly into the lair entrance or 1 drop/m in ants' paths for perimeter treatment.

Apply drops of gel having a diameter of about 1 cm (approximately 0.2 g)

Product can be used continuously for 2 - 3 months without replacing unconsumed gel. Maximum of 12 applications per year.

Risk mitigation measures: Avoid any unnecessary contact to the preparation. Misuse may cause health damage.

Do not contaminate food, feed, eating utensils or food contact surfaces.

Storage: Keep away from food, drink and animal feeding stuffs. Keep out from reach of children.

Precautions: After using IMIDASECT ANTS and before eating, wash your hands with soap and water. Do not apply the gel to surfaces where food is stored, prepared or consumed. Protect food, cooking utensils and cutlery from contamination. Do not apply the gel in places accessible to children and pets.

First aid:

IF ON SKIN: Wash with plenty of water.

If skin irritation or rash occurs: Get medical advice/attention.

24 h Emergency number: -NHS 111