



**RACAN  
DIFE BLOCK**  
Difenacoum  
(CAS 56073-07-5) : 0.0025%  
Warning: **H373 May cause damage  
to organs (blood) through  
prolonged or repeated exposure.**

P102: Keep out of reach of children.  
P103: Read label before use P220: Keep/  
store away from food, drink and animal  
feeding stuffs. P260: Avoid breathing  
dust. P262: Do not get on skin. P270: Do  
not eat, drink or smoke when using this  
product. P301+310: IF SWALLOWED:  
Immediately call a POISON CENTRE or  
doctor/physician. P314: Get medical  
advice/attention if you feel unwell.  
P404+405: Store locked up in a closed  
container. P501: Dispose of contents/  
container in accordance with applicable  
regulations.

**LODI UK - Building 69, Third Avenue,  
Pensnett Trading Estate, Kingswinford,  
West Midlands DY6 7FD, UK**  
sales@lodi-uk.com www.lodi-uk.com

#### ADVICE

Racan Dife Block is highly palatable and  
lethal to rats and mice. The active ingredient,  
difenacoum 25ppm, is a non-single feed  
poison available in the UK. This makes it  
ideal for use in and around the home.

Please note; a rat or a mouse requires  
multiple feeds of this bait to hit its lethal  
dose, so it is important to keep replenishing  
bait until the infestation is controlled.

**RACAN**  
DIFE BLOCK

L  
LodiUK

# RAT & MOUSE KILLER BLOCKBAIT



MULTI-FEED BAIT TECHNOLOGY



**300g**  
**10 x 30g  
BLOCKS**



**RACAN**  
DIFE BLOCK



**Home  
Defence**

MULTI-FEED BAIT  
TECHNOLOGY

Our **HOME DEFENCE** rodenticides  
have been carefully selected to give  
parents, young families and pet  
owners the confidence they have  
chosen a non-single feed poison  
effective enough to administer a  
lethal dose to rats and mice.

Specially designed, this multi-feed  
bait is consumed over a few days so  
rats and mice cannot associate the  
poison with making them feel unwell.  
Once the lethal dose is reached rats  
and mice will take several days for  
the anticoagulant to bring them to  
their untimely death.

RACAN Dife Blocks are supplied in  
special wrap. The unique wrapper  
allows the scents of the block to  
permeate and become palatable to  
mice and rats whilst ensuring you  
don't come into contact with the  
poison. Simply place the wrapped  
block into the RACAN bait station.

**This pack contains  
10 x 30g block bait**



0 700461 822899 >

**Authorisation No:** UK-2016-1053-0001

**User category:** For general public use.

**Field of use:** Indoor use for mice and rats / outdoor around  
buildings for rats only.

**Application method:** In tamper resistant bait stations.

**Target organism:** Rattus norvegicus (Brown rat) ; Mus musculus  
(House mouse).

**Application rate(s) and frequency:** For rats: 90-100g of bait per  
bait station. If more than one bait station is needed, the minimum  
distance between bait stations should be 5-10m apart. For mice:  
20-30g of bait per bait station. If more than 1 bait station is needed,  
the minimum distance between bait stations should be 3-5m apart.

**Use-specific instructions for use:** Mice: The bait stations  
should be visited at least every 2 to 3 days at the beginning of the  
treatment and at least weekly afterwards, in order to check whether  
the bait is accepted, the bait stations are intact and to remove  
rodent bodies. Re-fill bait when necessary. Rats:- Place the bait  
stations in areas non-labile to flooding. Replace any bait in a bait  
station in which bait has been damaged by water or contaminated  
by dirt. The bait stations should be visited only 5 to 7 days after the  
beginning of the treatment and at least weekly afterwards, in order  
to check whether the bait is accepted, the bait stations are intact  
and to remove rodent bodies. Refill bait when necessary.

**Instructions for use:** Read and follow the product information as  
well as any information accompanying the product or provided  
at the point of sale before using it. Prior to the use of rodenticide  
products, non-chemical control methods (e.g. traps) should be  
considered. Remove food which is readily attainable for rodents  
(e.g. spilled grain or food waste). Apart from this, do not clean up  
the infested area just before the treatment, as this only disturbs  
the rodent population and makes bait acceptance more difficult to  
achieve. Bait stations should be placed in the immediate vicinity  
where rodent activity has been observed (e.g. travel paths, nesting  
sites, feedlots, holes, burrows etc.). Where possible, bait stations  
must be fixed to the ground or other structures. Do not open the  
sachets containing the bait. Place bait stations out of the reach of  
children, birds, pets, farm animals and other non target  
animals. Place bait stations away from food, drink and animal  
feeding stuffs, as well as from utensils or surfaces that have  
contact with these. Do not place bait stations near water drainage  
systems where they can come into contact with water. When using  
the product do not eat, drink or smoke. Wash hands and directly  
exposed skin after using the product. Remove the remaining bait or  
the bait stations at the end of the treatment period.

**Risk mitigation measures:** Consider preventive control measures  
(plug holes, remove potential food and drinking as far as possible)  
to improve product intake and reduce the likelihood of re-invasion.  
Do not use anticoagulant rodenticides as permanent baits (e.g. for  
prevention of rodent infestation or to detect rodent activity). Using

**RACAN DIFE BLOCK**  
Block bait containing 0.0025% w/w of Difenacoum

**RACAN**  
DIFE BLOCK

this product should eliminate rodents within 35 days. The product  
information (i.e. label and/or leaflet) shall clearly recommend that in  
case of suspected lack of efficacy by the end of the treatment (i.e.  
rodent activity is still observed), the user should seek advice from  
the product supplier or call a pest control service. Search for and  
remove dead rodents during treatment, at least as often as baiting  
stations are inspected. Dispose dead rodents in accordance with  
local requirements. In the UK, poisoned rodents should be double-  
bagged using plastic bags and either disposed of in a household  
waste bin with a secure lid to prevent access of wildlife or pets or  
collected by a specialist waste contractor or the local authority.

**Particulars of likely direct or indirect effects, first aid  
instructions and emergency measures to protect the  
environment:** This product contains an anticoagulant substance.  
If ingested, symptoms, which may be delayed, may include  
nosebleed and bleeding gums. In severe cases, there may be  
bruising and blood present in the faeces or urine. Antidote: Vitamin  
K1 administered by medical/veterinary personnel only. In case of:  
Dermal exposure, wash skin with water and then with water and  
soap. Eye exposure, rinse eyes with eye-rinse liquid or water, keep  
eyes lids open at least 10 minutes. Oral exposure, rinse mouth  
carefully with water. Never give anything by mouth to unconscious  
person. Do not provoke vomiting. If swallowed, seek medical advice  
immediately and show the product's container or label. UK medical  
professionals should contact the National Poisons Information  
Service (www.npis.org) for further advice. Contact a veterinary  
surgeon in case of ingestion by a pet. Bait stations must be labelled  
with the following information: "do not move or open"; "contains  
a rodenticide"; "product name or authorisation number"; "active  
substance(s)" and "in case of accident, call a poison centre".  
Hazardous to wildlife.

**Instructions for safe disposal of the product and its packaging:**  
At the end of the treatment, dispose uneaten bait and the packaging  
in accordance with local requirements. In the UK waste bait should  
be double bagged in plastic bags and disposed of in a household  
waste bin with a secure lid to prevent access of wildlife or pets or  
taken to a civic amenity site. For information on civic amenity sites  
contact the local authority.

**Conditions of storage and shelf-life of the product under  
normal conditions of storage:** Store in a dry, cool and well  
ventilated place. Keep the container closed and away from direct  
sunlight. Store in places prevented from the access of children,  
birds, pets and farm animals. Shelf life: 2-years.

Because of their delayed mode of action, anticoagulant rodenticides  
take from 4 to 10 days to be effective after consumption of the bait.  
Rodents can be disease carriers. Do not touch dead rodents with  
bare hands, use gloves or use tools such as tongs when disposing  
them. This product contains a bittering agent and a dye.

Lot No. and Expiry Date: See pack.

L  
LodiUK

Marketed by Lodi UK Limited:  
Building 69, Third Avenue, Pensnett Trading Estate, Kingswinford,  
West Midlands DY6 7FD Tel: 01384 404242 Email: sales@lodi-uk.com  
www.lodi-uk.com

e 300g

LUK RACAN DIFE Block 300gr

E-4835°C